

From Pseudohypoparathyroidism to inactivating PTH/PTHrP Signalling Disorder
(iPPSD), a novel classification proposed by the European EuroPHP-network

Susanne Thiele^{1,*}, Giovanna Mantovani^{2,*}, Anne Barlier³, Valentina Boldrin²,
Paolo Bordogna², Luisa De Sanctis⁴, Francesca M Elli², Kathleen Freson⁵, Intza
Garin⁶, Virginie Grybek^{7,8}, Patrick Hanna^{7,8}, Benedetta Izzi⁵, Olaf Hiort¹, Beatriz
Lecumberri⁹, Arrate Pereda^{6,10}, Vrinda Saraff¹¹, Caroline Silve^{6,7,11}, Serap
Turan¹², Alessia Usardi⁶, Ralf Werner¹, Guiomar Perez de Nanclares^{5,*}, Agnès
Linglart^{6,7,14*}

¹Division of Experimental Pediatric Endocrinology and Diabetes, Department of
Pediatrics, University of Lübeck, Germany. ²Fondazione IRCCS Ca' Granda
Ospedale Maggiore Policlinico, Endocrinology and Diabetology Unit,
Department of Clinical Sciences and Community Health, University of Milan,
Milan, Italy. ³APHM, Hôpital la Conception, Laboratory of Molecular Biology,
13385, Marseille, France. ⁴Department of Public Health and Pediatric Sciences,
University of Torino, Italy. ⁵Department of Cardiovascular Sciences, Center for
Molecular and Vascular Biology, University of Leuven, Belgium. ⁶Molecular
(Epi)Genetics Laboratory, BioAraba National Health Institute, OSI Araba
University Hospital, Vitoria-Gasteiz, Spain. ⁷APHP, Reference Center for rare

disorders of the Calcium and Phosphate Metabolism, filière OSCAR and Plateforme d'Expertise Maladies Rares Paris-Sud, Hôpital Bicêtre Paris Sud, Le Kremlin Bicêtre, France. ⁸INSERM U1169, Hôpital Bicêtre, Le Kremlin Bicêtre, et Université Paris-Saclay, France. ⁹Department of Endocrinology and Nutrition, La Paz University Hospital, Madrid, Spain. ¹⁰Department of Biochemistry and Molecular Biology, University of Basque Country, Leioa, Spain. ¹¹Department of Endocrinology and Diabetes, Birmingham Children's Hospital, Birmingham, United Kingdom. ¹² APHP, Service de Biochimie et Génétique Moléculaires, Hôpital Cochin, Paris, France. ¹³Department of Pediatrics, Division of Endocrinology and Diabetes, Marmara University, Istanbul, Turkey. ¹⁴APHP, Department of paediatric endocrinology and diabetology, Bicêtre Paris Sud hospital, Le Kremlin Bicêtre, France.

*these authors equally contributed to the manuscript; all the other members of the EuroPHP are listed in alphabetical order.

Address all correspondence and requests for reprints to

agnes.linglart@aphp.fr

Key words: Pseudohypoparathyroidism, pseudopseudohypoparathyroidism, Gsa, *GNAS* mutation, *PRKAR1A* mutation, *PDE4D* mutation, *PDE3D* mutation,

42 *PTH1R* mutation, imprinting, inactivating PTH/PTHrP signalling disorders, new
43 classification

44

45 Word count: 3,866

46

47 **Abbreviated title:** iPPSD, a novel classification for PHP

48

49 **Abbreviations:** PHP: pseudohypoparathyroidism, PTH: parathyroid hormone,
50 PTH1R: PTH receptor type 1, PTHrP: PTH related peptide, Gsa: alpha subunit of
51 stimulatory G proteins, PHP1: PHP type 1, PHP2: PHP type 2, PHP1A: PHP type
52 1A, PHP1C: PHP type 1C, PHP1B: PHP type 1B, ACRDYS1: acrodysostosis type 1,
53 ACRDYS2: acrodysostosis type 2, AHO: Albright Hereditary Osteodystrophy,
54 PPHP: pseudopseudohypoparathyroidism, DMR: differentially methylated
55 regions, AD-PHP1B: autosomal dominant PHP1B, LOI: loss of imprinting, LOM:
56 loss of methylation, UPD: uniparental disomy PKA: protein kinase A, PDE:
57 phosphodiesterase, HTNB: Hypertension and Brachydactyly Syndrome, iPPSD:
58 Inactivating PTH/PTHrP Signalling Disorder, BDE: Brachydactyly type E, GHRH:
59 Growth Hormone Releasing Hormone, IUGR: Intra Uterine Growth Retardation,
60 ICR: Imprinting Control Region, POH: Progressive Osseous Heteroplasia

61

Abstract

Objective: Disorders caused by impairments in the parathyroid hormone (PTH) signalling pathway are historically classified under the term pseudohypoparathyroidism (PHP), that encompasses rare, related but highly heterogeneous diseases with demonstrated (epi)genetic causes. The actual classification is based on the presence or absence of specific clinical and biochemical signs together with an *in vivo* response to exogenous PTH and the results of an *in vitro* assay to measure Gsa protein activity. However, this classification disregards other related diseases like acrodysostosis (ACRDYS) or Progressive Osseous Heteroplasia (POH), as well as recent findings of clinical and genetic/epigenetic background of the different subtypes.

Therefore, the EuroPHP network decided to develop a new classification that encompasses all disorders with impairments in PTH and/or PTHrP cAMP-mediated pathway.

Design and Methods: Extensive review of the literature was performed. Several meetings were organised to discuss about a new, more effective and accurate way to describe disorders caused by abnormalities of the PTH/PTHrP signalling pathway.

82 **Results and Conclusions:** After determining the major and minor criteria
83 to be considered for the diagnosis of these disorders, we proposed to
84 group them under the term «inactivating PTH/PTHrP signalling
85 disorder», (iPPSD). This terminology: 1) defines the common mechanism
86 responsible for all diseases, 2) does not require a confirmed genetic
87 defect, 3) avoids ambiguous terms like “pseudo”, 4) eliminates the
88 clinical or molecular overlap between diseases. We believe that the use
89 of this nomenclature and classification will facilitate the development of
90 rationale and comprehensive international guidelines for the diagnosis
91 and treatment of iPPSDs.

92

Introduction

Pseudohypoparathyroidism (PHP) encompasses a group of rare, related, highly heterogeneous and deeply impairing disorders characterized by end-organ resistance to the action of parathyroid hormone (PTH) and in most instances associated with a demonstrated (epi)genetic component (1–3). PHP is historically the first hormone resistance syndrome described by Fuller Albright and colleagues in 1942 (4).

A better understanding of the PHP pathophysiology followed the identification of the PTH receptor (PTH1R) and its signal transduction pathway (Figure 1) (5,6). PTH1R, through its activation by two ligands, the PTH and the PTH related peptide (PTHrP), regulates skeletal development, bone turnover and mineral ion homeostasis. In the kidney, binding of PTH to PTH1R stimulates the production of 1,25 dihydroxy vitamin D3, and inhibits phosphate reabsorption in the proximal tubule, while it increases calcium reabsorption in the distal nephron. In the growth plate, PTHrP promotes endochondral ossification, through the binding to PTH1R (7).

112 The Blomstrand chondrodysplasia (OMIM # 215045), a lethal form of dwarfism
113 (8), was the first disorder associated with biallelic loss-of-function mutations of
114 the *PTH1R* gene (9). Subsequently, one report has described a milder
115 phenotype in living children affected with Eiken disease (OMIM # 600002),
116 short stature, elevated PTH and mutations of *PTH1R* (10,11).

117 A defect in the response of the proximal renal tubule to PTH is the hallmark of
118 all forms of PHP. It manifests as hypocalcemia, hyperphosphatemia and
119 elevated circulating levels of PTH in the absence of vitamin D deficiency
120 (5,7,12).

121 PTH receptor couples with the stimulatory G protein (Gsa), leading to cAMP
122 formation. Renal tubular response to exogenously administered PTH through
123 measurement of serum and urinary cAMP levels, permits the differentiation of
124 PHP type 1 (PHP1), in which a blunted cAMP response is observed, from PHP
125 type 2 (PHP2), where cAMP increase is conserved but the phosphaturic
126 response is deficient (13). To date, only a handful of PHP2 cases have been
127 reported, and the molecular defect responsible for this variant is still unknown.
128 It has also been hypothesized that PHP2 could either be an acquired defect
129 secondary to vitamin D deficiency (14), as calcium and vitamin D
130 supplementation resulted in normalization of the phosphaturic response to

PTH in some patients (14,15) or due to defects downstream the Gsa protein, as seen in patients with acrodysostosis type 1 (ACRDYS1) (16).

In 1980, deficiency in the Gsa protein activity in erythrocytes extracted from patients affected with PHP1 was demonstrated *in vitro* (17,18). For years, this bioassay allowed the diagnosis of PHP, and contributed to PHP subclassification (see below).

PHP type 1 (PHP1) is further subdivided based on the presence (PHP1A and PHP1C; OMIM #103580 and #612462, respectively) (6,17–19) or absence (PHP1B; OMIM #603233) (6,20) of Albright Hereditary Osteodystrophy (AHO) (Table 1). AHO is a clinical entity initially described together with PHP in 1942, which encompasses heterogeneous clinical findings such as brachydactyly, rounded face, short stature, stocky build and subcutaneous ossifications (4,21,22). Additional features that may not directly relate to AHO, yet extensively associated with PHP1A individuals, include obesity, varying degrees of intellectual disability and resistance to several hormones, including TSH, GHRH and calcitonin (23–28). The subcategory of PHP1C has all the characteristics of PHP1A, except that Gsa activity in erythrocytes was found comparable to controls (29,30).

151 Interestingly, patients showing the physical features of AHO without any
152 evidence of PTH resistance were also described by Albright and colleagues ten
153 years after their first report of PHP (21). This new syndrome, named pseudo-
154 pseudohypoparathyroidism (PPHP; OMIM #612463) may be present either in
155 kindreds with PHP or as an isolated defect. It is possible that the “bone
156 phenotype” observed in AHO is largely mediated by the resistance to PTHrP at
157 the growth plate during fetal and post-natal growth (31).

158
159 In 1990 the first heterozygous inactivating mutation in the gene coding for Gsa
160 (*GNAS*), responsible for PHP1A, was described (32). Since then, several Gsa-
161 coding mutations have been identified in all of its 13 exons with different
162 frequency, with a detection rate of about 70% (33–39). Cases of deletions of
163 20q, including part or the whole *GNAS* gene, and an inversion at *GNAS* have
164 been recently reported (40–44). Remarkably, similar mutations when
165 paternally inherited, or occurring *de novo* on the paternal allele of *GNAS* may
166 lead to PPHP or to Progressive Osseous Heteroplasia (POH, OMIM #166350), a
167 disorder characterized by heterotopic ossifications expanding into deep
168 muscles and connective tissues (45,46).

170 *GNAS* is a locus encoding several transcripts through alternative splicing. In
171 most tissues, except for *Gsa*, the *GNAS* transcripts are of monoallelic origin due
172 to the control of their expression by parent-specific differentially methylated
173 regions (DMRs) (Figure 2) (47). In thyroid, pituitary gland and most likely in the
174 proximal tubule (36), *Gsa* is predominantly expressed from the maternal allele
175 through a yet unexplained mechanism (48,49). In the early 2000, the molecular
176 defect of PHP1B was characterized. The most consistent defect common to all
177 PHP1B patients is a paternal-specific pattern of cytosine methylation within the
178 maternal *GNAS A/B*: transcriptional start site (TSS)-DMR (*GNAS A/B*:TSS-DMR;
179 previously known as exon *A/B* or 1A) which could lead to a decreased
180 expression of *Gsa* in the renal proximal tubules, hence PTH resistance (50).
181 Fifteen to 20% of the PHP1B cases present familial history with an autosomal
182 dominant mode of inheritance (AD-PHP1B) through the maternal lineage. Most
183 AD-PHP1B show loss of imprinting (LOI) limited to the *GNAS A/B*:TSS-DMR
184 (more precisely a loss of methylation [LOM]) associated with deletions on the
185 maternal allele of *cis*-acting control elements within *STX16* or *NESP55* (51–55),
186 although other maternally inherited deletions have been identified affecting all
187 four DMRs (*GNAS-NESP*:TSS-DMR, *GNAS-AS1*:TSS-DMR, *GNAS-XL*:Ex1-DMR and
188 *GNAS A/B*:TSS-DMR) (56–58).

189 The remaining cases of PHP1B are sporadic. They present with broad LOI at
190 *GNAS*, including the *GNAS A/B:TSS-DMR*. The molecular basis of this broad LOI
191 is yet to be identified, with an exception of less than 10% of the patients who
192 are affected by paternal complete or segmental uniparental disomy (UPD) of
193 the chromosome 20, comprising the *GNAS* locus (59–63).

194

195 To summarize, the existing classification of PHP (Table 1) is based on the
196 following criteria, 1-presence or absence of AHO differentiates **PHP1A/PHP1C**
197 from **PHP1B**, 2-presence or absence of hormonal resistance differentiates **PHP1**
198 from **PPHP**, 3-*in vivo* response to exogenous PTH as for nephrogenic cAMP
199 synthesis and phosphaturia separates **PHP1** from **PHP2**, and 4-*in vitro* assay
200 measuring the Gsa protein activity from erythrocyte membranes, differentiates
201 between **PHP1A** and **PHP1C**.

202

203 As described above, the existing PHP classification does not include molecular
204 defect as a criterion and fails to stratify PHP and AHO as well as include
205 conditions such as acrodysostosis, POH and PTH1R-related chondrodysplasia. In
206 this manuscript, we therefore propose to review the rationale of this
207 nomenclature and recommend a novel classification for disorders impairing the
208 PTH/PTHrP signalling pathway.

Methodology

The EuroPHP network met on three different occasions (October 2014, May 2015, November 2015) to discuss and agree on a novel classification. The aims of these meetings were 1- to identify the limitations in the current PHP classification, 2- to formulate mandatory criteria for the new classification, 3- to propose a comprehensive definition gathering all the disorders, 4- to analyse the classifications used in other genetic/epigenetic conditions, 5- to generate a novel classification. The methodology comprised of a thorough review of the current literature to facilitate comparison and form basis for the proposal of a new classification.

We have carefully considered a series of classifications proposed for various rare genetic/epigenetic disorders, including the reporting manuscripts that were taken into consideration for the design of a novel classification (summarized in Table 2). In brief, methodologies were similar. A group of experts in the field identified the deficiencies in the existing classification/terminology and the need for an update. Subsequently agreement on a novel terminology and classification was reached and reported (64–70).

229 Challenges and limitations of the current classification

230 Recent clinical and molecular data gathered for these complex disorders have
231 questioned the distinction of the different PHP and AHO subtypes in the
232 existing classification (Table 1). We have selected the following limits of the
233 current classification.

234 **1-** In a subset of patients with PHP1A and varying degree of AHO, LOI of *GNAS*
235 identical to that of PHP1B has been reported, suggesting a molecular and
236 clinical overlap between the two subtypes (71); further confirmed (72–75).

237 **2-** PHP1B patients present with a moderate reduction of Gsa activity in
238 erythrocyte membranes, reminiscent -yet less severe- to that observed in
239 patients with PHP1A and PPHP (76).

240 **3-** Recently, mild resistance to PTH was described in patients affected with
241 PPHP, carrying a paternal *GNAS* mutation (77), showing that the hormonal
242 resistance is not only associated with maternally inherited *GNAS* mutations.

243 **4-** Different molecular defects have been identified in patients with PHP1C, i.e.
244 LOI at *GNAS* and four loss-of-function mutations in the *GNAS* carboxyl-terminus
245 leading to a conserved adenylyl cyclase receptor-independent activation but
246 disrupted receptor-mediated activation (29,30,78).

247 **6-** Paternal *GNAS* mutations associated with Progressive Osseous Heteroplasia
248 are usually truncating mutations (79), yet they are identical to those found in

249 families with PHP1A and/or PPHP (45). Also noteworthy, is that a fraction of
250 POH patients exhibits some of the typical AHO features and, conversely, some
251 PHP1A patients carrying mutations on the maternal allele, present with
252 progressive deepening heterotopic ossifications. The hypothesis that POH
253 should be considered as a form of PPHP is therefore debated (80,81).

254 **7-** Heterozygous mutations in *PRKAR1A* -coding for the regulatory subunit of
255 the protein kinase A (PKA)- and *PDE4D* -coding for phosphodiesterase type 4-
256 have been found in patients with acrodysostosis (16,82,83). Acrodysostosis
257 refers to a heterogeneous group of rare diseases characterized by skeletal
258 dysplasia and characteristic features, including brachydactyly, facial
259 dysmorphism and, in some cases, mental retardation (84–88). Acrodysostosis
260 differs from PHP by more generalized osseous abnormalities (87,89).
261 Resistance to PTH and/or TSH is present in about 60-70% of acrodysostosis
262 patients with a *PRKAR1A* mutation, while, in case of a *PDE4D* mutation, such
263 hormone resistances are found only in a smaller subset of 10-20%.
264 Interestingly, few patients bearing a *PRKAR1A* mutation have been described in
265 patients with a phenotype indistinguishable from PHP1A (90,91).

266 **8-** Heterozygous mutations in *PDE3A* have been identified in patients affected
267 with hypertension and brachydactyly type E (Hypertension and Brachydactyly
268 Syndrome (HTNB): OMIM #112410) (92).

269 9- Disorders associated with an impaired function of *PTH1R*, i.e. the Blomstrand
270 and Eiken skeletal dysplasia, are currently not included in the classification.

271

272 Over the past two decades it became obvious that clinical features such as AHO
273 or *in vitro* assays such as Gsa bioactivity fail to differentiate between PHP
274 subtypes. In addition, mutations of genes different from *GNAS* have been
275 shown to lead to PTH and/or PTHrP resistance and *GNAS* mutations might
276 trigger diseases different from PHP/PPHP (i.e. POH). These disorders are not
277 encompassed by the current classification system.

278

279 For all these reasons, different independent studies from the authors of the
280 present paper, as well as the “EuroPHP network” concluded and agreed that a
281 uniform terminology is required to create a functional working classification
282 that reflects the current knowledge of the diseases (29,93,94).

283

284 **Terminology**

285 We propose the term of «inactivating PTH/PTHrP signalling disorder»,
286 abbreviated as iPPSD, which encompass all disorders related to this pathway.

287 We also propose that numbering will allow for both clinical features as well as
288 molecular and genetic findings to be included. The advantages of this
289 terminology are as follows: 1- it describes the common mechanism responsible

for the diseases, 2- it does not require a confirmed genetic defect, 3- it avoids the ambiguous term like “pseudo”, 4- it eliminates the clinical or molecular overlap between diseases and 5- it is flexible to incorporate new evolving information.

We recognize that the nomenclature «inactivating PTH/PTHrP signalling disorder» might be initially difficult for patients and caregivers to remember. It would therefore be helpful to rely on the abbreviation iPPSD. Equally the former terms “pseudohypoparathyroidism” and “pseudopseudohypoparathyroidism” were also long and challenging to use for communication. PTH/PTHrP specific pathway was deliberately included in the name of the classification to avoid the misperception with disorders resulting from the inactivation of G protein-coupled receptors, i.e. inactivating mutations in the TSH receptor or in the FSH receptor. All nomenclature based on the cAMP signalling were carefully considered and rejected due to their generic nature.

Identification of mandatory criteria for the new classification

Basis for the newly proposed classification of iPPSD are:

- To provide patients with an unambiguous diagnosis;

- 310 • To base nomenclature on pathophysiology, i.e. the
311 PTH1R/Gsa/cAMP/PKA pathway, and a standardized diagnostic
312 pathway;
- 313 • To formulate basis to develop new therapeutic approaches;
- 314 • To be sufficiently flexible and adaptable to include emerging clinical
315 and molecular information;
- 316 • To be simple and usable for the caregivers.

317

318 It is therefore of significant importance to define the category of iPPSD a
319 patient belongs to, based on the characterization of clinical/biochemical
320 criteria, to facilitate a definitive diagnosis and, if possible, through molecular
321 analysis, a more specific denomination within the classification.

322

323 We suggest three key clinical features as major criteria for the diagnosis of
324 iPPSD. The proposed major criteria have minimum or no overlap with other
325 conditions due to different mechanisms (see Table 3, especially for the
326 differential diagnoses).

327 We also propose a list of minor criteria that are associated with iPPSD. These
328 are less specific to iPPSD compared to major criteria and can occur in other

clinical conditions. Therefore, minor criteria need to be combined with one or more major criteria to establish the diagnosis of iPPSD.

Major criteria

1. PTH resistance

The hallmark of inactivating PTH/PTHrP signalling disorders is the resistance of the renal proximal tubule to the action of PTH. All genetic defects leading to a deficient PTH1R signalling in the kidney will therefore be named iPPSD.

2. Ectopic ossifications

Ectopic ossifications are superficial, subcutaneous nodules, defined as ectopic bone formation in the adipose or dermal tissue. Progressive Osseous Heterotopic calcifications often begin in the dermal and subcutaneous tissues and later progress to the deeper tissues, such as muscles and tendons. In children, ectopic ossifications are highly suggestive of an inactivating *GNAS* mutation, i.e. iPPSD.

3. Brachydactyly

Brachydactyly refers to shortening of fingers, toes or both. Brachydactyly type E (BDE, OMIM #113300) encompasses variable shortening of the metacarpals/metatarsals, often with the involvement of phalanges (see Figure 3). It can either present in isolation or as part of a genetic disorder, most of which are included among iPPSD (95).

349 Brachydactyly can be challenging to identify in early childhood, and tends to
350 become more evident during early puberty. Brachydactyly can be overlooked
351 when all bones are short as in acrodysostosis since early childhood.

352 While PTH resistance and ectopic ossifications are considered major criteria for
353 iPPSD, brachydactyly is less specific and should therefore be combined with at
354 least one major or two minor criteria to trigger the diagnosis of iPPSD.

355 **Minor criteria**

356 *1. Thyroid Stimulating Hormone (TSH) resistance*

357 In iPPSD, TSH resistance is often mild and characterized by elevated TSH levels
358 associated with free thyroxine (T4) levels in a normal or low-normal reference
359 range. This occurs in the absence of goitre and markers of autoimmune disease
360 (26,27). TSH resistance can sometimes be the first detected sign of iPPSD,
361 especially in countries where screening for congenital hypothyroidism is
362 routinely performed (96).

363 *2. Other hormone resistances*

364 Very few other hormone resistances have been demonstrated so far.
365 Resistance to growth hormone releasing hormone (GHRH), leading to growth
366 hormone deficiency, is the most frequent additional resistance found in PHP1A,
367 affecting as many as 60% of patients (97–99). Calcitonin resistance has been
368 described without clinical features in patients affected with PHP1A (27).

Elevated follicular stimulating hormone (FSH) and luteinizing hormone (LH) levels were reported both by us and Namnoum et al (78,100). Glucagon and epinephrine resistances were demonstrated in patients with features of PHP and low Gsa bioactivity through *in vivo* testing (6,101).

3. Motor and cognitive retardation or impairment

Psychomotor and cognitive alterations have been described in about 40 to 70% of the patients with a maternal coding mutation of *GNAS* (25,34), as well as in some patients affected with acrodysostosis (83,85,86). Psychiatric manifestations have also been reported in these patients (102). Patients with paternal mutations of *GNAS* or epigenetic modifications of the *GNAS* DMRs seem unaffected (25,63).

4. Intra-uterine and post-natal growth retardation

Intra-uterine growth retardation (IUGR) has been frequently observed in both maternal and paternal inherited inactivating *GNAS* coding mutations. However, IUGR is more pronounced in patients harbouring mutations on the paternal *GNAS* allele, mainly when affecting *GNAS* exon 2-13 mutations, compared to patients with *GNAS* exon 1/intron 1 mutations (103). IUGR has also been described in acrodysostosis with mutations in *PRKAR1A* or *PDE4D*, and in patients with mutations in *PDE3A* (16,82,90,92). A LOI at the maternal *GNAS* A/B: TSS-DMR has been associated with increased intra-uterine growth (104).

389 Post-natal growth retardation is a frequent sign in PHP1A and acrodysostosis.
390 Growth hormone deficiency and premature closure of the epiphysis result in
391 short stature (16,82,97,105). Growth retardation has also been observed in
392 PHP1B, although only in exceptional cases (71,74), and in patients with Eiken
393 dysplasia (10).

394 5. *Obesity/overweight*

395 Obesity or overweight may be the most nonspecific minor sign, however it
396 occurs very frequently in disorders with an impaired PTH/PTHrP signalling
397 pathway and may help to differentiate between the different subtypes. Growth
398 hormone deficiency, impaired lipolytic response of epinephrine (101), or
399 decreased resting energy expenditure (106) contribute to the development of
400 obesity in patients with mutations on the maternal allele of *GNAS* (23,107).
401 Obesity is also a frequent feature in patients affected with acrodysostosis
402 (16,90,108).

403 6. *Flat nasal bridge and/or maxillar hypoplasia and/or round face*

404 Elements of facial dysmorphism have been associated with acrodysostosis (flat
405 nasal bridge and/or maxillar hypoplasia) or with PHP1A (round face) (4,86).

406

407 **Diagnosis of iPPSD**

We propose that a minimum of one of the major criteria is mandatory for the clinical diagnosis of iPPSD. PTH resistance or ectopic ossifications may lead to the diagnosis of iPPSD with or without the presence of minor criteria. However, brachydactyly type E (BDE) should be associated with at least one major or two minor criteria to suggest iPPSD, as it is a common feature of several other diseases and syndromes (Table 3).

The known molecular causes of PTH/PTHrP signalling disorders are:

- Inactivating mutations of *PTH1R*;
- Inactivating heterozygous mutations in the coding sequence of *GNAS*-Gsa;
- Methylation changes of the DMRs of *GNAS* caused by
 - deletions or duplications at ICRs (*STX16*; *NESP*; *GNAS-AS1*);
 - paternal UPD of chromosome 20q;
 - unknown mechanism(s);
- Heterozygous mutations of *PRKAR1A*;
- Heterozygous mutations of *PDE4D*;
- Heterozygous mutations of *PDE3A*.

In contrast to the former diagnostic classification based solely on the phenotype, once iPPSD has been identified (using criteria described Table 3),

428 we propose to further subtype iPPSD based on the underlying molecular
429 (epi)genetic defect. Therefore, the term iPPSD will refer to the pathophysiology
430 of the PTH/PTHrP signalling abnormalities, while the number will refer to the
431 underlying molecular mechanism (responsible for the pathology). We have
432 numbered iPPSD subtypes starting with *PTH1R* mutations.

433

434 **The novel classification of iPPSD**

435 The European PHP-network proposes the following classification (Figure 4):

- 436 • iPPSD: clinical/biochemical diagnosis based on the major/minor criteria
437 as defined above, in the absence of genetic investigation;
- 438 • iPPSD1: loss of function mutation in *PTH1R* ;
- 439 • iPPSD2: loss of function mutation in *Gsa*;
- 440 • iPPSD3: methylation change(s) at one or more *GNAS* DMRs, associated
441 with or without a genetic (deletion) or cytogenetic (UPD) defect;
- 442 • iPPSD4: *PRKAR1A* mutation;
- 443 • iPPSD5: *PDE4D* mutation;
- 444 • iPPSD6: *PDE3A* mutation;
- 445 • iPPSDx: lack of genetic/epigenetic defect identified following molecular
446 investigation of known genes described above;

- iPPSDn+1: the identification of a novel gene/molecular defect will lead to a disease named iPPSD7, then 8 and so on.

iPPSD3 encompasses all disorders associated with changes in the methylation patterns of the DMRs of *GNAS*, including UPD(20)pat and deletion within *STX16*, *NESP*, etc. Of most significance, is the common mechanism shared by these patients, i.e. the LOM at the *GNAS* A/B:TSS-DMR. Grouping them under iPPSD3 highlights this common mechanism. Secondly, we anticipated the difficulties in integrating the multiplicity of the epigenetic mechanisms within the classification system as this adds no further diagnostic value. However, the further specification of the epigenetic defect can remain part of a private exchange between the molecular laboratory, the patient and his/her physician. We recommend the use of Arabic numerals to avoid the confusion with letters (II with the number 11 for example).

The advantages of this new nomenclature are: 1- it stratifies the disorders into clusters caused by the same mechanism, 2- it is flexible and open to accommodate new defects to be discovered in the future and 3- it simplifies the concept of the overlapping disorders under a single umbrella.

467 This classification however bears some limitations. We deliberately did not
468 include the parental origin of the genetic/epigenetic defect, although some
469 iPPSD are imprinting disorders -namely iPPSD2 and iPPSD3- and their
470 phenotypic expression depends on their parental inheritance. The main reason
471 behind this being the association of PTH resistance and POH with both
472 maternal and paternal inactivating *GNAS* mutations. Therefore, the mechanism
473 of the two allelic *GNAS* mutations can be considered alike. However, in daily
474 practice, the parental origin of the *GNAS* defect should be considered,
475 particularly for genetic counselling. In fact, AHO and multiple hormone
476 resistance including PTH resistance are largely associated with maternal *GNAS*
477 coding defects, whereas, isolated AHO and/or POH are more often associated
478 with paternal *GNAS* coding defects.

479 Another limitation of this classification is the inability to sub classify individuals
480 with a pure clinical suspicion of iPPSD and lack of complete (epi)genetic testing.
481 While such patients cannot be classified as iPPSDx or with a specific number,
482 we recommend that they are classified as iPPSD.

483 The inclusion of the disorders involving the two main ligands of the PTH1R, i.e.
484 hypoparathyroidism (109) and brachydactyly type E with short stature
485 (mutations in *PTH1R* the gene encoding PTHrP (110,111)) to the classification
486 may be argued. However, we decided to exclude them due to several other

487 issues such as, 1- their different biochemical pattern including low levels of PTH
488 responsible for hypoparathyroidism 2- the dramatic difference in the therapy
489 of hypoparathyroidism and defects in PTH1R signalling, respectively and 3- the
490 difference in research goals in the two disease groups.

491

492 **Perspectives**

493 We believe that the use of the new nomenclature will facilitate a more
494 straightforward approach to the diagnosis of iPPSD, increase awareness of the
495 red-flag signs of PTH resistance, ectopic ossifications and brachydactyly type E.
496 It would allow for the classification of patients into local catalogues used by the
497 different healthcare organizations in a more homogenous way, and enable
498 future observational and research studies in the field.

499

500 We strongly believe that too many denominations for similar diseases and
501 patients with phenocopies (PHP, PPHP, POH, ACRDYS, TRPS, BDE, AHO) has
502 diluted and dispersed research advance, adding undue complexity to the
503 causative mechanism and proved challenging for the experts in building a
504 global research network in the field.

505

506 Regular use of the classification in daily practice or for scientific purposes will
507 allow appropriate amendments in the best interest of the patients.

508

509 While producing this novel nomenclature and classification, we have identified
510 the need to 1) disseminate this alternative classification to be positively
511 enriched by the clinical and scientific community, 2) validate the major/minor

criteria in a series of patients affected by different iPPSDs, and 3) develop international guidelines for the diagnosis and treatment of the iPPSDs in the near future.

Declaration of interest

All the authors declare that there is no conflict of interest that could be perceived as prejudicing the impartiality of the research reported.

Funding

This work and/or members of the network Euro-Pseudohypoparathyroidism network (EuroPHP) was supported by grants obtained from the European Society for Paediatric Endocrinology (ESPE) Research Unit, the University of Luebeck (H02-2011), the Italian Ministry of Health [GR-2009-1608394], Ricerca Corrente Funds [to Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico], the Instituto de Salud Carlos III and European fund for Regional Development, FEDER, [PI13/00467 to GPdN], the Basque Department of Health [GV2014111017 to GPdN], the University of Basque Country (Ref: 48198) [to AP], the I3SNS Program of the Spanish Ministry of Health [CP03/0064; SIVI 1395/09 to GPdN], the APHP Bicêtre Paris-Sud Reference Center for Rare

531 Disorders of the Calcium and Phosphate Metabolism [to AL, CS, VG, PH and AU]
532 as well as the Plateforme d'Expertise Maladies Rares Paris-Sud.

533

534 All members of the EuroPHP network are members of the EUCID.net (COST
535 action BM1208 on imprinting disorders; www.imprinting-disorders.eu), which
536 was supported by SANDOZ FRANCE for training actions.

537

538

539 **Legends for figures**

540

541 **Figure 1:** Schematic transduction of PTH1R/Gsa/cAMP/PKA pathway.

542 Upon ligand binding (PTH or PTHrP are mentioned on the figure), the receptor

543 (PTH1R) activates the G protein. Then the Gsa subunit triggers the activation of

544 the adenylate cyclase leading to cAMP synthesis. cAMP binds to the regulatory

545 1A subunits (R1A) of the PKA, the most common effector of cAMP.

546 Upon cAMP binding, the catalytic subunits (Cat) dissociate from the R1A

547 subunits, and phosphorylate numerous target proteins including CREB (cAMP-

548 responsive binding elements) and the phosphodiesterases (PDEs). CREB

549 activates the transcription of cAMP responsive genes. Intracellular cAMP is

550 then deactivated by PDEs, among which are PDE4D and PDE3A.

551 PTH1R: transmembrane convolutional black line; G protein: trimer α , β , γ ;

552 cAMP: grey diamond; PKA: tetramer R1A (regulatory subunit 1A) and Cat

553 (catalytic subunit); phosphodiesterases: ovals PDE4D or PDE3A; DNA: scale bar.

554

555 **Figure 2:** The imprinted human *GNAS* locus (Hg19-chr20:57,414,795-556 57,486,250), on chromosome 20, close to the *STX16* gene (Hg19-

557 chr20:57,226,309-57,254,5812) (source UCSC, Hg19). The

558 centromeric/telomeric orientation of the chromosome is indicated. The

559 maternal (NESP), paternal (AB, AS and XL) and biallelic (Gsa) transcripts are
560 depicted as arrows. Maternal and paternal expressed transcripts are drawn
561 above and below the horizontal line, respectively. Black boxes: coding exons;
562 grey boxes: non-coding exons; arrows: transcription (direction and parental
563 origin). The brackets delimit the imprinting control element deletions, which
564 have been reported. *STX16* gene: full brackets: the recurrent *STX16* deletion of
565 3.3kb (38); large dotted brackets: the *STX16* deletion of 4.4 kb (39); small
566 dotted brackets: the *STX16* deletion of 29,5kb (42). *GNAS* locus: full brackets:
567 the 4.7kb and 4kb deletions removing the *NESP* exon and exons 3 and 4 of
568 *GNAS-AS1* (40); large dotted brackets: the 4.2kb deletion removing exons 3 and
569 4 of *GNAS-AS1* (43); deletions of 40pb [*] and 33pb [#] in introns of *NESP* and
570 *GNAS-AS1* (44); small dotted brackets: the *NESP* and *GNAS-AS1* deletion (41).
571 Mat: maternal ; Pat: paternal ; cen: centromeric ; tel: telomeric.

572

573 **Figure 3:** Patterns of brachydactyly type E associated with iPPSD.

574 **a-e**, brachydactylies associated with coding mutations in the Gsa subunit of the
575 G protein (iPPSD2). **f-g**, bone phenotype associated with the loss of imprinting
576 at the *GNAS* locus (iPPSD3). **h-j**, brachydactylies associated with the molecular
577 defect in *PRKAR1A* (iPPSD4) and *PDE4D* (iPPSD5). Note the phenotypic overlap
578 between a, h, j, and b, c, g, i, respectively.

579

580 **Figure 4:** Schematic representation of the new classification proposed by the
581 European PHP network.

582 According to the suggested new classification Blomstrand and Eiken
583 chondrodysplasia, PHP type 1 & 2, PPHP, AHO, POH and acrodysostosis
584 clinically/biochemically diagnosed without genetic investigation are named
585 iPPSD; Blomstrand and Eiken chondrodysplasia due to PTHR1 inactivating
586 mutations are named iPPSD1; PHP1A, PHP1C, PPHP and POH clinically
587 diagnosed and characterized by Gsa inactivating mutations are termed iPPSD2;
588 PHP1B clinically diagnosed and due to methylation changes at the *GNAS* DMRs
589 is classified as iPPSD3; in the presence of acrodysostosis type 1 or *PRKAR1A*
590 mutations, the disease is classified as iPPSD4. Acrodysostosis type 2 or *PDE4D*
591 mutations are termed iPPSD5; *PDE3A* mutations are categorized as iPPSD6;
592 patients lacking genetic or epigenetic defects at the known genes fall under the
593 category of iPPSDx; any newly discovered genetic/molecular defects will be
594 labelled as iPPSDn+1.

595

596

597 **References**

- 598 1. Mantovani G. Clinical review: Pseudohypoparathyroidism: diagnosis and
599 treatment. *J Clin Endocrinol Metab.* oct 2011;96(10):3020-30.
- 600 2. Weinstein LS, Yu S, Warner DR, Liu J. Endocrine manifestations of
601 stimulatory G protein alpha-subunit mutations and the role of genomic
602 imprinting. *Endocr Rev.* oct 2001;22(5):675-705.
- 603 3. Turan S, Bastepe M. The GNAS complex locus and human diseases
604 associated with loss-of-function mutations or epimutations within this
605 imprinted gene. *Horm Res Paediatr.* 2013;80(4):229-41.
- 606 4. Albright F, Burnett CH, Smith PH, Parson W. Pseudohypoparathyroidism -
607 an example of « Seabright-Bantam syndrome ». *Endocrinology.*
608 1942;30:922-32.
- 609 5. Chase LR, Melson GL, Aurbach GD. Pseudohypoparathyroidism: defective
610 excretion of 3',5'-AMP in response to parathyroid hormone. *J Clin Invest.*
611 1969;48:1832-44.
- 612 6. Levine MA, Downs RW Jr, Moses AM, Breslau NA, Marx SJ, Lasker RD, et
613 al. Resistance to multiple hormones in patients with
614 pseudohypoparathyroidism. Association with deficient activity of guanine
615 nucleotide regulatory protein. *Am J Med.* 1983;74:545-56.
- 616 7. Cheloha RW, Gellman SH, Vilardaga J-P, Gardella TJ. PTH receptor-1
617 signalling-mechanistic insights and therapeutic prospects. *Nat Rev*
618 *Endocrinol.* déc 2015;11(12):712-24.
- 619 8. Blomstrand S, Claesson I, Säve-Söderbergh J. A case of lethal congenital
620 dwarfism with accelerated skeletal maturation. *Pediatr Radiol.*
621 1985;15:141-3.
- 622 9. Zhang P, Jobert AS, Couvineau A, Silve C. A homozygous inactivating
623 mutation in the parathyroid hormone/parathyroid hormone-related
624 peptide receptor causing Blomstrand chondrodysplasia. *J Clin Endocrinol*
625 *Metab.* 1998;83:3365-8.
- 626 10. Duchatelet S, Ostergaard E, Cortes D, Lemainque A, Julier C. Recessive
627 mutations in PTHR1 cause contrasting skeletal dysplasias in Eiken and
628 Blomstrand syndromes. *Hum Mol Genet.* 1 janv 2005;14(1):1-5.

- 629 11. Hoogendam J, Farih-Sips H, Wÿnaendts LC, Löwik CWGM, Wit JM,
630 Karperien M. Novel mutations in the parathyroid hormone (PTH)/PTH-
631 related peptide receptor type 1 causing Blomstrand
632 osteochondrodysplasia types I and II. *J Clin Endocrinol Metab.* mars
633 2007;92(3):1088-95.
- 634 12. Stone M, Hosking D, Garcia-Himmelstine C, White D, Rosenblum D, Worth
635 H. The renal response to exogenous parathyroid hormone in treated
636 pseudohypoparathyroidism. *Bone.* 1993;14:727-35.
- 637 13. Drezner M, Neelon FA, Lebovitz HE. Pseudohypoparathyroidism type II: a
638 possible defect in the reception of the cyclic AMP signal. *N Engl J Med.* 15
639 nov 1973;289(20):1056-60.
- 640 14. Rao DS, Parfitt AM, Kleerekoper M, Pumo BS, Frame B. Dissociation
641 between the effects of endogenous parathyroid hormone on adenosine
642 3',5'-monophosphate generation and phosphate reabsorption in
643 hypocalcemia due to vitamin D depletion: an acquired disorder
644 resembling pseudohypoparathyroidism type II. *J Clin Endocrinol Metab.*
645 août 1985;61(2):285-90.
- 646 15. Rodriguez HJ, Villarreal H Jr, Klahr S, Slatopolsky E.
647 Pseudohypoparathyroidism type II: restoration of normal renal
648 responsiveness to parathyroid hormone by calcium administration. *J Clin*
649 *Endocrinol Metab.* oct 1974;39(4):693-701.
- 650 16. Linglart A, Menguy C, Couvineau A, Auzan C, Gunes Y, Cancel M, et al.
651 Recurrent PRKAR1A mutation in acrodysostosis with hormone resistance.
652 *N Engl J Med.* 9 juin 2011;364(23):2218-26.
- 653 17. Levine MA, Downs RW Jr, Singer M, Marx SJ, Aurbach GD, Spiegel AM.
654 Deficient activity of guanine nucleotide regulatory protein in erythrocytes
655 from patients with pseudohypoparathyroidism. *Biochem Biophys Res*
656 *Commun.* 1980;94:1319-24.
- 657 18. Farfel Z, Brickman AS, Kaslow HR, Brothers VM, Bourne HR. Defect of
658 receptor-cyclase coupling protein in psudohypoparathyroidism. *N Engl J*
659 *Med.* 31 juill 1980;303(5):237-42.
- 660 19. Radeke HH, Auf'mkolk B, Jüppner H, Krohn HP, Keck E, Hesch RD. Multiple
661 pre- and postreceptor defects in pseudohypoparathyroidism (a

- 662 multicenter study with twenty four patients). J Clin Endocrinol Metab.
663 1986;62:393-402.
- 664 20. Silve C, Santora A, Breslau N, Moses A, Spiegel A. Selective resistance to
665 parathyroid hormone in cultured skin fibroblasts from patients with
666 pseudohypoparathyroidism type 1b. J Clin Endocrinol Metab.
667 1986;62:640-4.
- 668 21. Albright F, Forbes AP, Henneman PH. Pseudo-pseudohypoparathyroidism.
669 Trans Assoc Am Physicians. 1952;65:337-50.
- 670 22. Eyre WG, Reed WB. Albright's hereditary osteodystrophy with cutaneous
671 bone formation. Arch Dermatol. déc 1971;104(6):634-42.
- 672 23. Long DN, McGuire S, Levine MA, Weinstein LS, Germain-Lee EL. Body
673 mass index differences in pseudohypoparathyroidism type 1a versus
674 pseudopseudohypoparathyroidism may implicate paternal imprinting of
675 Galpha(s) in the development of human obesity. J Clin Endocrinol Metab.
676 mars 2007;92(3):1073-9.
- 677 24. Farfel Z, Friedman E. Mental deficiency in pseudohypoparathyroidism
678 type I is associated with Ns-protein deficiency. Ann Intern Med. août
679 1986;105(2):197-9.
- 680 25. Mouallem M, Shaharabany M, Weintrob N, Shalitin S, Nagelberg N,
681 Shapira H, et al. Cognitive impairment is prevalent in
682 pseudohypoparathyroidism type 1a, but not in
683 pseudopseudohypoparathyroidism: possible cerebral imprinting of
684 Galpha. Clin Endocrinol (Oxf). févr 2008;68(2):233-9.
- 685 26. Balavoine AS, Ladsous M, Velayoudom FL, Vlaeminck V, Cardot-Bauters C,
686 d'Herbomez M, et al. Hypothyroidism in patients with
687 pseudohypoparathyroidism type 1a: clinical evidence of resistance to TSH
688 and TRH. Eur J Endocrinol. oct 2008;159(4):431-7.
- 689 27. Vlaeminck-Guillem V, D'Herbomez M, Pigny P, Fayard A, Bauters C,
690 Decoulx M, et al. Pseudohypoparathyroidism 1a and hypercalcitoninemia.
691 J Clin Endocrinol Metab. juill 2001;86(7):3091-6.
- 692 28. Linglart A, Maupetit-Méhouas S, Silve C. GNAS -Related Loss-of-Function
693 Disorders and the Role of Imprinting. Horm Res Paediatr. 29 mars
694 2013;119-29.

- 695 29. Brix B, Werner R, Staedt P, Struve D, Hiort O, Thiele S. Different pattern of
696 epigenetic changes of the GNAS gene locus in patients with
697 pseudohypoparathyroidism type 1c confirm the heterogeneity of
698 underlying pathomechanisms in this subgroup of
699 pseudohypoparathyroidism and the demand for a new classification of
700 GNAS-related disorders. J Clin Endocrinol Metab. août
701 2014;99(8):E1564-70.
- 702 30. Thiele S, de Sanctis L, Werner R, Grotzinger J, Aydin C, Juppner H, et al.
703 Functional characterization of GNAS mutations found in patients with
704 pseudohypoparathyroidism type 1c defines a new subgroup of
705 pseudohypoparathyroidism affecting selectively Gsalpha-receptor
706 interaction. Hum Mutat. juin 2011;32(6):653-60.
- 707 31. Hirai T, Chagin AS, Kobayashi T, Mackem S, Kronenberg HM. Parathyroid
708 hormone/parathyroid hormone-related protein receptor signaling is
709 required for maintenance of the growth plate in postnatal life. Proc Natl
710 Acad Sci U S A. 4 janv 2011;108(1):191-6.
- 711 32. Patten JL, Johns DR, Valle D, Eil C, Gruppuso PA, Steele G, et al. Mutation
712 in the gene encoding the stimulatory G protein of adenylate cyclase in
713 Albright's hereditary osteodystrophy. New Engl J Med. 1990;322:1412-9.
- 714 33. Elli FM, deSanctis L, Ceoloni B, Barbieri AM, Bordogna P, Beck-Peccoz P, et
715 al. Pseudohypoparathyroidism type 1a and pseudo-
716 pseudohypoparathyroidism: the growing spectrum of GNAS inactivating
717 mutations. Hum Mutat. mars 2013;34(3):411-6.
- 718 34. Thiele S, Werner R, Grötzinger J, Brix B, Staedt P, Struve D, et al. A positive
719 genotype-phenotype correlation in a large cohort of patients with
720 Pseudohypoparathyroidism Type 1a and Pseudo-
721 pseudohypoparathyroidism and 33 newly identified mutations in the
722 GNAS gene. Mol Genet Genomic Med. mars 2015;3(2):111-20.
- 723 35. Lemos MC, Thakker RV. GNAS mutations in Pseudohypoparathyroidism
724 type 1a and related disorders. Hum Mutat. janv 2015;36(1):11-9.
- 725 36. Weinstein LS, Gejman PV, Friedman E, Kadowaki T, Collins RM, Gershon
726 ES, et al. Mutations of the Gs alpha-subunit gene in Albright hereditary
727 osteodystrophy detected by denaturing gradient gel electrophoresis. Proc
728 Natl Acad Sci U A. nov 1990;87(21):8287-90.

- 729 37. Mantovani G, Bondioni S, Linglart A, Maghnie M, Cisternino M, Corbetta
730 S, et al. Genetic analysis and evaluation of resistance to thyrotropin and
731 growth hormone-releasing hormone in pseudohypoparathyroidism type
732 1b. *J Clin Endocrinol Metab.* sept 2007;92(9):3738-42.
- 733 38. Ham H-J, Baek K-H, Lee J-Y, Kim SY, Mo EY, Kim ES, et al. Analysis of
734 aberrantly spliced transcripts of a novel de novo GNAS mutant in a male
735 with albright hereditary osteodystrophy and PHP1A. *Horm Metab Res*
736 *Horm Stoffwechselforschung Horm Métabolisme.* juill 2015;47(8):585-90.
- 737 39. Lin MH, Numbenjapon N, Germain-Lee EL, Pitukcheewanont P.
738 Progressive osseous heteroplasia, as an isolated entity or overlapping
739 with Albright hereditary osteodystrophy. *J Pediatr Endocrinol Metab*
740 *JPEM.* juill 2015;28(7-8):911-8.
- 741 40. Geneviève D, Sanlaville D, Faivre L, Kottler M-L, Jambou M, Gosset P, et al.
742 Paternal deletion of the GNAS imprinted locus (including Gnasxl) in two
743 girls presenting with severe pre- and post-natal growth retardation and
744 intractable feeding difficulties. *Eur J Hum Genet EJHG.* sept
745 2005;13(9):1033-9.
- 746 41. Fernandez-Rebollo E, García-Cuartero B, Garin I, Largo C, Martínez F,
747 Garcia-Lacalle C, et al. Intragenic GNAS deletion involving exon A/B in
748 pseudohypoparathyroidism type 1A resulting in an apparent loss of exon
749 A/B methylation: potential for misdiagnosis of
750 pseudohypoparathyroidism type 1B. *J Clin Endocrinol Metab.* févr
751 2010;95(2):765-71.
- 752 42. Fernandez-Rebollo E, Barrio R, Pérez-Nanclares G, Carcavilla A, Garin I,
753 Castaño L, et al. New mutation type in pseudohypoparathyroidism type
754 1a. *Clin Endocrinol (Oxf).* nov 2008;69(5):705-12.
- 755 43. Mitsui T, Nagasaki K, Takagi M, Narumi S, Ishii T, Hasegawa T. A family of
756 pseudohypoparathyroidism type 1a with an 850-kb submicroscopic
757 deletion encompassing the whole GNAS locus. *Am J Med Genet A.* janv
758 2012;158A(1):261-4.
- 759 44. Garin I, Elli FM, Linglart A, Silve C, de Sanctis L, Bordogna P, et al. Novel
760 microdeletions affecting the GNAS locus in pseudohypoparathyroidism:
761 characterization of the underlying mechanisms. *J Clin Endocrinol Metab.*
762 avr 2015;100(4):E681-7.

- 763 45. Adegbite NS, Xu M, Kaplan FS, Shore EM, Pignolo RJ. Diagnostic and
764 mutational spectrum of progressive osseous heteroplasia (POH) and other
765 forms of GNAS-based heterotopic ossification. *Am J Med Genet A*. 15 juill
766 2008;146A(14):1788-96.
- 767 46. Shore EM, Ahn J, Jan de Beur S, Li M, Xu M, Gardner RJM, et al. Paternally
768 inherited inactivating mutations of the GNAS1 gene in progressive
769 osseous heteroplasia. *N Engl J Med*. 10 janv 2002;346(2):99-106.
- 770 47. Hayward B, Bonthron D. An imprinted antisense transcript at the human
771 GNAS1 locus. *Hum Mol Genet*. 2000;9:835-41.
- 772 48. Yu S, Yu D, Lee E, Eckhaus M, Lee R, Corria Z, et al. Variable and tissue-
773 specific hormone resistance in heterotrimeric Gs protein alpha-subunit
774 (Gsalph) knockout mice is due to tissue-specific imprinting of the gsalph
775 gene. *Proc Natl Acad Sci U S A*. 21 juill 1998;95(15):8715-20.
- 776 49. Mantovani G, Ballare E, Giammona E, Beck-Peccoz P, Spada A. The
777 gsalph gene: predominant maternal origin of transcription in human
778 thyroid gland and gonads. *J Clin Endocrinol Metab*. oct
779 2002;87(10):4736-40.
- 780 50. Liu J, Litman D, Rosenberg MJ, Yu S, Biesecker LG, Weinstein LS. A GNAS1
781 imprinting defect in pseudohypoparathyroidism type 1b. *J Clin Invest*. nov
782 2000;106(9):1167-74.
- 783 51. Bastepe M, Fröhlich LF, Hendy GN, Indridason OS, Josse RG, Koshiyama H,
784 et al. Autosomal dominant pseudohypoparathyroidism type 1b is
785 associated with a heterozygous microdeletion that likely disrupts a
786 putative imprinting control element of GNAS. *J Clin Invest*. oct
787 2003;112(8):1255-63.
- 788 52. Linglart A, Gensure RC, Olney RC, Jüppner H, Bastepe M. A novel STX16
789 deletion in autosomal dominant pseudohypoparathyroidism type 1b
790 redefines the boundaries of a cis-acting imprinting control element of
791 GNAS. *Am J Hum Genet*. mai 2005;76(5):804-14.
- 792 53. Bastepe M, Fröhlich LF, Linglart A, Abu-Zahra HS, Tojo K, Ward LM, et al.
793 Deletion of the NESP55 differentially methylated region causes loss of
794 maternal GNAS imprints and pseudohypoparathyroidism type 1b. *Nat*
795 *Genet*. janv 2005;37(1):25-7.

- 796 54. Richard N, Abeguilé G, Coudray N, Mittre H, Gruchy N, Andrieux J, et al. A
797 new deletion ablating NESP55 causes loss of maternal imprint of A/B
798 GNAS and autosomal dominant pseudohypoparathyroidism type 1b. *J Clin*
799 *Endocrinol Metab.* mai 2012;97(5):E863-7.
- 800 55. Elli FM, de Sanctis L, Peverelli E, Bordogna P, Pivetta B, Miolo G, et al.
801 Autosomal dominant pseudohypoparathyroidism type 1b: a novel
802 inherited deletion ablating STX16 causes loss of imprinting at the A/B
803 DMR. *J Clin Endocrinol Metab.* avr 2014;99(4):E724-8.
- 804 56. Chillambhi S, Turan S, Hwang D-Y, Chen H-C, Jüppner H, Bastepe M.
805 Deletion of the noncoding GNAS antisense transcript causes
806 pseudohypoparathyroidism type 1b and biparental defects of GNAS
807 methylation in cis. *J Clin Endocrinol Metab.* août 2010;95(8):3993-4002.
- 808 57. Rezwan FI, Poole RL, Prescott T, Walker JM, Karen Temple I, Mackay DJG.
809 Very small deletions within the NESP55 gene in
810 pseudohypoparathyroidism type 1b. *Eur J Hum Genet EJHG.* avr
811 2015;23(4):494-9.
- 812 58. Takatani R, Molinaro A, Grigelioniene G, Tafaj O, Watanabe T, Reyes M, et
813 al. Analysis of Multiple Families with Single Individuals Affected by
814 Pseudohypoparathyroidism Type 1b (PHP1B) Reveals Only One Novel
815 Maternally Inherited GNAS Deletion. *J Bone Miner Res Off J Am Soc Bone*
816 *Miner Res.* 19 oct 2015;
- 817 59. Bastepe M, Lane AH, Jüppner H. Paternal uniparental isodisomy of
818 chromosome 20q--and the resulting changes in GNAS1 methylation--as a
819 plausible cause of pseudohypoparathyroidism. *Am J Hum Genet.* mai
820 2001;68(5):1283-9.
- 821 60. Fernández-Rebollo E, Lecumberri B, Garin I, Arroyo J, Bernal-Chico A, Goñi
822 F, et al. New mechanisms involved in paternal 20q disomy associated with
823 pseudohypoparathyroidism. *Eur J Endocrinol Eur Fed Endocr Soc.* déc
824 2010;163(6):953-62.
- 825 61. Dixit A, Chandler KE, Lever M, Poole RL, Bullman H, Mughal MZ, et al.
826 Pseudohypoparathyroidism type 1b due to paternal uniparental disomy of
827 chromosome 20q. *J Clin Endocrinol Metab.* janv 2013;98(1):E103-8.
- 828 62. Takatani R, Minagawa M, Molinaro A, Reyes M, Kinoshita K, Takatani T, et
829 al. Similar frequency of paternal uniparental disomy involving

- 830 chromosome 20q (patUPD20q) in Japanese and Caucasian patients
831 affected by sporadic pseudohypoparathyroidism type 1b (sporPHP1B).
832 Bone. oct 2015;79:15-20.
- 833 63. Maupetit-Méhouas S, Azzi S, Steunou V, Sakakini N, Silve C, Reynes C, et
834 al. Simultaneous hyper- and hypomethylation at imprinted loci in a subset
835 of patients with GNAS epimutations underlies a complex and different
836 mechanism of multilocus methylation defect in
837 pseudohypoparathyroidism type 1b. Hum Mutat. août
838 2013;34(8):1172-80.
- 839 64. Van Dijk FS, Pals G, Van Rijn RR, Nikkels PGJ, Cobben JM. Classification of
840 Osteogenesis Imperfecta revisited. Eur J Med Genet. févr 2010;53(1):1-5.
- 841 65. June RR, Aggarwal R. The use and abuse of diagnostic/classification
842 criteria. Best Pract Res Clin Rheumatol. déc 2014;28(6):921-34.
- 843 66. Bonafe L, Cormier-Daire V, Hall C, Lachman R, Mortier G, Mundlos S, et al.
844 Nosology and classification of genetic skeletal disorders: 2015 revision.
845 Am J Med Genet A. 23 sept 2015;
- 846 67. Eckardt K-U, Alper SL, Antignac C, Bleyer AJ, Chauveau D, Dahan K, et al.
847 Autosomal dominant tubulointerstitial kidney disease: diagnosis,
848 classification, and management-A KDIGO consensus report. Kidney Int.
849 oct 2015;88(4):676-83.
- 850 68. Marcucci G, Cianferotti L, Beck-Peccoz P, Capezzone M, Cetani F, Colao A,
851 et al. Rare diseases in clinical endocrinology: a taxonomic classification
852 system. J Endocrinol Invest. févr 2015;38(2):193-259.
- 853 69. Picard C, Al-Herz W, Bousfiha A, Casanova J-L, Chatila T, Conley ME, et al.
854 Primary Immunodeficiency Diseases: an Update on the Classification from
855 the International Union of Immunological Societies Expert Committee for
856 Primary Immunodeficiency 2015. J Clin Immunol. 19 oct 2015;
- 857 70. Alberti KG, Zimmet PZ. Definition, diagnosis and classification of diabetes
858 mellitus and its complications. Part 1: diagnosis and classification of
859 diabetes mellitus provisional report of a WHO consultation. Diabet Med J
860 Br Diabet Assoc. juill 1998;15(7):539-53.
- 861 71. de Nanclares GP, Fernandez-Rebollo E, Santin I, Garcia-Cuartero B,
862 Gaztambide S, Menendez E, et al. Epigenetic defects of GNAS in patients
863 with pseudohypoparathyroidism and mild features of Albright's

- 864 hereditary osteodystrophy. J Clin Endocrinol Metab. juin
865 2007;92(6):2370-3.
- 866 72. Unluturk U, Harmanci A, Babaoglu M, Yasar U, Varli K, Bastepe M, et al.
867 Molecular diagnosis and clinical characterization of
868 pseudohypoparathyroidism type-Ib in a patient with mild Albright's
869 hereditary osteodystrophy-like features, epileptic seizures, and defective
870 renal handling of uric acid. Am J Med Sci. juill 2008;336(1):84-90.
- 871 73. Mariot V, Maupetit-Méhouas S, Sinding C, Kottler M-L, Linglart A. A
872 maternal epimutation of GNAS leads to Albright osteodystrophy and
873 parathyroid hormone resistance. J Clin Endocrinol Metab. mars
874 2008;93(3):661-5.
- 875 74. Mantovani G, de Sanctis L, Barbieri AM, Elli FM, Bollati V, Vaira V, et al.
876 Pseudohypoparathyroidism and GNAS epigenetic defects: clinical
877 evaluation of albright hereditary osteodystrophy and molecular analysis
878 in 40 patients. J Clin Endocrinol Metab. févr 2010;95(2):651-8.
- 879 75. Izzi B, Decallonne B, Devriendt K, Bouillon R, Vanderschueren D,
880 Levtchenko E, et al. A new approach to imprinting mutation detection in
881 GNAS by Sequenom EpiTYPER system. Clin Chim Acta Int J Clin Chem. 14
882 déc 2010;411(23-24):2033-9.
- 883 76. Zazo C, Thiele S, Martín C, Fernandez-Rebollo E, Martinez-Indart L,
884 Werner R, et al. Gs α activity is reduced in erythrocyte membranes of
885 patients with psedohypoparathyroidism due to epigenetic alterations at
886 the GNAS locus. J Bone Miner Res. août 2011;26(8):1864-70.
- 887 77. Turan S, Thiele S, Tafaj O, Brix B, Atay Z, Abali S, et al. Evidence of
888 hormone resistance in a pseudo-pseudohypoparathyroidism patient with
889 a novel paternal mutation in GNAS. Bone. févr 2015;71:53-7.
- 890 78. Linglart A, Carel JC, Garabédian M, Lé T, Mallet E, Kottler ML. GNAS1
891 lesions in pseudohypoparathyroidism Ia and Ic: genotype phenotype
892 relationship and evidence of the maternal transmission of the hormonal
893 resistance. J Clin Endocrinol Metab. janv 2002;87(1):189-97.
- 894 79. Lebrun M, Richard N, Abeguilé G, David A, Coëslier Dieux A, Journal H, et
895 al. Progressive osseous heteroplasia: a model for the imprinting effects of
896 GNAS inactivating mutations in humans. J Clin Endocrinol Metab. juin
897 2010;95(6):3028-38.

- 898 80. Kaplan FS, Shore EM. Progressive osseous heteroplasia. J Bone Min Res.
899 2000;15(11):2084-94.
- 900 81. Eddy MC, De Beur SM, Yandow SM, McAlister WH, Shore EM, Kaplan FS,
901 et al. Deficiency of the alpha-subunit of the stimulatory G protein and
902 severe extraskeletal ossification. J Bone Min Res. 2000;15(11):2074-83.
- 903 82. Michot C, Le Goff C, Goldenberg A, Abhyankar A, Klein C, Kinning E, et al.
904 Exome sequencing identifies PDE4D mutations as another cause of
905 acrodysostosis. Am J Hum Genet. 6 avr 2012;90(4):740-5.
- 906 83. Lee H, Graham JM Jr, Rimoin DL, Lachman RS, Krejci P, Thompson SW, et al.
907 Exome sequencing identifies PDE4D mutations in acrodysostosis. Am J
908 Hum Genet. 6 avr 2012;90(4):746-51.
- 909 84. Maroteaux P, Malamut G. [Acrodysostosis]. Presse Médicale. 27 nov
910 1968;76(46):2189-92.
- 911 85. Robinow M, Pfeiffer RA, Gorlin RJ, McKusick VA, Renuart AW, Johnson GF,
912 et al. Acrodysostosis. A syndrome of peripheral dysostosis, nasal
913 hypoplasia, and mental retardation. Am J Dis Child 1960. mars
914 1971;121(3):195-203.
- 915 86. Reiter S. Acrodysostosis. A case of peripheral dysostosis, nasal hypoplasia,
916 mental retardation and impaired hearing. Pediatr Radiol. 10 avr
917 1978;7(1):53-5.
- 918 87. Davies SJ, Hughes HE. Familial acrodysostosis: can it be distinguished from
919 Albright's hereditary osteodystrophy? Clin Dysmorphol. oct
920 1992;1(4):207-15.
- 921 88. Silve C, Le-Stunff C, Motte E, Gunes Y, Linglart A, Clauser E. Acrodysostosis
922 syndromes. BoneKEy Rep. 2012;1:225.
- 923 89. Ablow RC, Hsia YE, Brandt IK. Acrodysostosis coinciding with
924 pseudohypoparathyroidism and pseudo-pseudohypoparathyroidism. AJR
925 Am J Roentgenol. janv 1977;128(1):95-9.
- 926 90. Linglart A, Fryssira H, Hiort O, Holterhus P-M, Perez de Nanclares G,
927 Argente J, et al. PRKAR1A and PDE4D mutations cause acrodysostosis but
928 two distinct syndromes with or without GPCR-signaling hormone
929 resistance. J Clin Endocrinol Metab. déc 2012;97(12):E2328-38.

- 930 91. Nagasaki K, Iida T, Sato H, Ogawa Y, Kikuchi T, Saitoh A, et al. PRKAR1A
931 mutation affecting cAMP-mediated G protein-coupled receptor signaling
932 in a patient with acrodysostosis and hormone resistance. *J Clin Endocrinol*
933 *Metab.* sept 2012;97(9):E1808-13.
- 934 92. Maass PG, Aydin A, Luft FC, Schächterle C, Weise A, Stricker S, et al.
935 PDE3A mutations cause autosomal dominant hypertension with
936 brachydactyly. *Nat Genet.* juin 2015;47(6):647-53.
- 937 93. Mantovani G, Elli FM, Spada A. GNAS epigenetic defects and
938 pseudohypoparathyroidism: time for a new classification? *Horm Metab*
939 *Res Horm Stoffwechselforschung Horm Métabolisme.* sept
940 2012;44(10):716-23.
- 941 94. Bastepe M. Genetics and epigenetics of parathyroid hormone resistance.
942 *Endocr Dev.* 2013;24:11-24.
- 943 95. Pereda A, Garin I, Garcia-Barcina M, Gener B, Beristain E, Ibañez AM, et al.
944 Brachydactyly E: isolated or as a feature of a syndrome. *Orphanet J Rare*
945 *Dis.* 12 sept 2013;8:141.
- 946 96. Romanet P, Osei L, Netchine I, Pertuit M, Enjalbert A, Reynaud R, et al.
947 Case report of GNAS epigenetic defect revealed by a congenital
948 hypothyroidism. *Pediatrics.* avr 2015;135(4):e1079-83.
- 949 97. Germain-Lee EL, Groman J, Crane JL, Jan de Beur SM, Levine MA. Growth
950 hormone deficiency in pseudohypoparathyroidism type 1a: another
951 manifestation of multihormone resistance. *J Clin Endocrinol Metab.* sept
952 2003;88(9):4059-69.
- 953 98. Mantovani G, Maghnie M, Weber G, De Menis E, Brunelli V, Cappa M, et
954 al. Growth hormone-releasing hormone resistance in
955 pseudohypoparathyroidism type 1a: new evidence for imprinting of the Gs
956 alpha gene. *J Clin Endocrinol Metab.* sept 2003;88(9):4070-4.
- 957 99. de Sanctis L, Bellone J, Salerno M, Faleschini E, Caruso-Nicoletti M,
958 Cicchetti M, et al. GH secretion in a cohort of children with
959 pseudohypoparathyroidism type 1a. *J Endocrinol Invest.* févr
960 2007;30(2):97-103.
- 961 100. Namnoum AB, Merriam GR, Moses AM, Levine MA. Reproductive
962 dysfunction in women with Albright's hereditary osteodystrophy. *J Clin*
963 *Endocrinol Metab.* 1998;83:824-9.

- 964 101. Carel JC, Le Stunff C, Condamine L, Mallet E, Chaussain JL, Adnot P, et al.
 965 Resistance to the lipolytic action of epinephrine: a new feature of protein
 966 Gs deficiency. *J Clin Endocrinol Metab.* nov 1999;84(11):4127-31.
- 967 102. Kadilli I, Colicchio S, Guglielmo R, Vollono C, Della Marca G, Janiri L.
 968 Clinical insights by the presence of bipolar disorder in
 969 pseudohypoparathyroidism type 1A. *Gen Hosp Psychiatry.* oct
 970 2015;37(5):497.e3-5.
- 971 103. Richard N, Molin A, Coudray N, Rault-Guillaume P, Jüppner H, Kottler M-L.
 972 Paternal GNAS mutations lead to severe intrauterine growth retardation
 973 (IUGR) and provide evidence for a role of XL α s in fetal development. *J Clin*
 974 *Endocrinol Metab.* sept 2013;98(9):E1549-56.
- 975 104. Bréhin A-C, Colson C, Maupetit-Méhouas S, Grybek V, Richard N, Linglart
 976 A, et al. Loss of methylation at GNAS exon A/B is associated with
 977 increased intrauterine growth. *J Clin Endocrinol Metab.* avr
 978 2015;100(4):E623-31.
- 979 105. Germain-Lee EL. Short stature, obesity, and growth hormone deficiency in
 980 pseudohypoparathyroidism type 1a. *Pediatr Endocrinol Rev PER.* avr
 981 2006;3 Suppl 2:318-27.
- 982 106. Shoemaker AH, Lomenick JP, Saville BR, Wang W, Buchowski MS, Cone
 983 RD. Energy expenditure in obese children with
 984 pseudohypoparathyroidism type 1a. *Int J Obes* 2005. août
 985 2013;37(8):1147-53.
- 986 107. Roizen JD, Danzig J, Groleau V, McCormack S, Casella A, Harrington J, et
 987 al. Resting Energy Expenditure is Decreased in Pseudohypoparathyroidism
 988 Type 1A. *J Clin Endocrinol Metab.* 28 déc 2015;jc20153895.
- 989 108. Lynch DC, Dymont DA, Huang L, Nikkel SM, Lacombe D, Campeau PM, et
 990 al. Identification of novel mutations confirms PDE4D as a major gene
 991 causing acrodysostosis. *Hum Mutat.* janv 2013;34(1):97-102.
- 992 109. Bilezikian JP, Khan A, Potts JT, Brandi ML, Clarke BL, Shoback D, et al.
 993 Hypoparathyroidism in the adult: epidemiology, diagnosis,
 994 pathophysiology, target-organ involvement, treatment, and challenges
 995 for future research. *J Bone Miner Res Off J Am Soc Bone Miner Res.* oct
 996 2011;26(10):2317-37.

- 997 110. Klopocki E, Hennig BP, Dathe K, Koll R, de Ravel T, Baten E, et al. Deletion
998 and point mutations of PTHLH cause brachydactyly type E. *Am J Hum*
999 *Genet.* 12 mars 2010;86(3):434-9.
- 1000 111. Thomas-Teinturier C, Pereda A, Garin I, Diez-Lopez I, Linglart A, Silve C, et
1001 al. Report of two novel mutations in PTHLH associated with brachydactyly
1002 type E and literature review. *Am J Med Genet A.* 6 déc 2015;
- 1003 112. Schwindinger WF, Miric A, Zimmerman D, Levine MA. A novel Gs alpha
1004 mutant in a patient with Albright hereditary osteodystrophy uncouples
1005 cell surface receptors from adenylyl cyclase. *J Biol Chem.* 14 oct
1006 1994;269(41):25387-91.
- 1007 113. Srivastava T, Krudys J, Mardis NJ, Sebestyen-VanSickle J, Alon US.
1008 Cinacalcet as adjunctive therapy in pseudohypoparathyroidism type 1b.
1009 *Pediatr Nephrol Berl Ger.* 1 déc 2015;
- 1010 114. Bakker B, Sonneveld LJJ, Woltering MC, Bikker H, Kant SG. A Girl With
1011 Beckwith-Wiedemann Syndrome and Pseudohypoparathyroidism Type 1B
1012 Due to Multiple Imprinting Defects. *J Clin Endocrinol Metab.* nov
1013 2015;100(11):3963-6.
- 1014 115. Perez-Nanclares G, Velayos T, Vela A, Muñoz-Torres M, Castaño L.
1015 Pseudohypoparathyroidism type 1b associated with novel duplications in
1016 the GNAS locus. *PloS One.* 2015;10(2):e0117691.
- 1017 116. Mehraein Y, Pfob M, Steinlein O, Aichinger E, Eggert M, Bubendorff V, et
1018 al. 2q37.3 Deletion Syndrome: Two Cases with Highly Distinctive Facial
1019 Phenotype, Discordant Association with Schizophrenic Psychosis, and
1020 Shared Deletion Breakpoint Region on 2q37.3. *Cytogenet Genome Res.*
1021 2015;146(1):33-8.
- 1022 117. Lee H, Graham Jr. JM, Rimoin DL, Lachman RS, Krejci P, Thompson SW, et
1023 al. Exome Sequencing Identifies PDE4D Mutations in Acrodysostosis. *Am J*
1024 *Hum Genet.* 6 avr 2012;90(4):746-51.
- 1025 118. Muhn F, Klopocki E, Graul-Neumann L, Uhrig S, Colley A, Castori M, et al.
1026 Novel mutations of the PRKAR1A gene in patients with acrodysostosis.
1027 *Clin Genet.* déc 2013;84(6):531-8.
- 1028 119. Kaname T, Ki C-S, Niikawa N, Baillie GS, Day JP, Yamamura K-I, et al.
1029 Heterozygous mutations in cyclic AMP phosphodiesterase-4D (PDE4D)

- 1030 and protein kinase A (PKA) provide new insights into the molecular
1031 pathology of acrodysostosis. *Cell Signal.* nov 2014;26(11):2446-59.
- 1032 120. Lindstrand A, Grigelioniene G, Nilsson D, Pettersson M, Hofmeister W,
1033 Anderlid B-M, et al. Different mutations in PDE4D associated with
1034 developmental disorders with mirror phenotypes. *J Med Genet.* janv
1035 2014;51(1):45-54.
- 1036 121. Mitsui T, Kim O-H, Hall CM, Offiah A, Johnson D, Jin D-K, et al.
1037 Acroscyphodysplasia as a phenotypic variation of
1038 pseudohypoparathyroidism and acrodysostosis type 2. *Am J Med Genet*
1039 *A.* oct 2014;164A(10):2529-34.
- 1040

Table 1: Former classification of PHP along with the other disorders affecting the PTH/PTHrP signalling pathway; note the overlap of phenotypes and molecular defects of the patients. Diseases included in the former classification are PHP1A, PHP1B, PHP1C and PPHP.

| | PHP1A | PHP1C | PHP1B | | | | | |
|--------------------------|---|--|--|--|--|---|----------------|--|
| Clinical presentation | AHO | AHO | No AHO | AHO in some patients (brachydactyly, subcutaneous ossification) and/or obesity | AHO in very few patients | Mental retardation reported in 2 patients, lambdoid synostosis, early-onset obesity, macrocephaly | BWS | No AHO |
| | Obesity | Obesity | No obesity | | Obesity may be present | | Obesity | no obesity |
| | Cognitive impairment | Cognitive impairment | No Cognitive impairment | | | | | |
| | Subcutaneous ossifications | Subcutaneous ossifications | | | | | | |
| Hormone resistance | Resistance to PTH, TSH, GHRH, calcitonin, epinephrine, glucagon and gonadotropins | Resistance to PTH, TSH, epinephrine and gonadotropins | PTH resistance | PTH resistance, ± TSH resistance | PTH resistance, ± TSH resistance | PTH resistance, ± TSH resistance | PTH resistance | PTH resistance, ± TSH resistance |
| In vitro activity of Gsa | Significantly below controls | Similar to controls | Similar to controls | Mildly decreased when compared to controls | Mildly decreased when compared to controls | | | Similar to controls |
| LOI at the GNAS DMRs | | | | LOM at the GNAS A/B:TSS-DMR | Broad LOI | Broad LOI | Broad LOI | Broad LOI |
| Genetic lesion | Heterozygous mutation in the coding sequence of GNAS (maternal allele) | Heterozygous mutation in the coding sequence of GNAS (p.E392K, p.E392X, p.L388R and p.Y391X, all in exon 13) (maternal allele) | Heterozygous mutation in the coding sequence of GNAS (p.Ile382del) (maternal allele) | Recurrent 3-kb STX16 deletion or 4.2-kb deletion of STX16 | unknown | UPD(20)pat including GNAS | MLID | Maternal deletion of NESP and/or AS or Duplication of GNAS |
| References | (32–39) | (29,30,78) | (112) | (50–52,113) | (50,51,63,74) | (59–63) | (114) | (53,54,58,115) |

| | PPHP | | POH | 2q37.3 Deletion Syndrome | PHP2 | Acrodysostosis | | | Blomstrand dysplasia | Eiken disease |
|---|---|---|--|--|---------------------------------------|---|--|---|--|--|
| Clinical presentation | AHO | AHO | Subcutaneous ossifications | AHO | No AHO | Severe AHO | AHO | Severe AHO | Lethal dwarfism | Epiphyseal dysplasia |
| | Subcutaneous ossifications | Subcutaneo us ossifications | | Cognitive impairment | Hypocalc emia, osteomal acia | Cognitive impairment in some patients | | Hypertensi on | | Short stature |
| Hormone resistance | No | Mild | No | No | PTH resistance | PTH resistance, and TSH in some patients | PTH resistance, and TSH in some patients | No | | Elevated PTH in one patient |
| In vitro activity of Gsa | Significantly below controls | Significantly below controls | | | | | | | | |
| LOI at the GNAS DMRs | | | | | | | | | | |
| Genetic lesion | Heterozygous mutation in the coding sequence of GNAS (paternal allele) | Heterozygou s mutation in the coding sequence of GNAS (paternal allele) | Heterozygous mutation in the coding sequence of GNAS (paternal allele) or no mutation identified | Deletion of the 2q37.3 chromosomal region including HDAC4 | None | Heterozygous mutation in the coding sequence of PRKAR1A or PDE4D | Heterozygo us mutation in the coding sequence of PRKAR1A | Heterozyg ous mutation in the coding sequence of PDE3A | Biallelic inactivating mutation in the coding sequence of PTH1R | Biallelic inactivating mutation in the coding sequence of PTH1R |
| References | (34,36,44,100) | (77) | (39,45,46,79,80) | (116) | (14) | (16,82,90,91,117-120) | (90,91) | (92) | (9,11) | (10) |

Legend: PHP, pseudohypoparathyroidism; PPHP, Pseudopseudohypoparathyroidism; AHO, Albright's hereditary osteodystrophy; BWS: Beckwith-Wiedemann syndrome; MLID: multilocus imprinting defect; NA, not available.

Table 2: Non exhaustive review of classifications used in other conditions

| | Methodology used to build the classification | Mode of classification | Advantages | Limitations |
|---|--|---|--|--|
| Primary Immunodeficiency Diseases (69) | 2-days meeting | Groups of diseases according to the most fundamental defect Presented as a table format | Allows a practical clinical framework for PID diagnosis | The complexities of these conditions cannot easily be captured in the limited table format |
| Skeletal dysplasia (66) | Meeting, extensive review of the literature, and circulation of drafts of the manuscript | Groups of diseases defined by molecular, biochemical and/or radiographic criteria | Disorders are caused by disturbances in related metabolic pathways or gene networks, Sheer number of conditions included | The “hybrid” nature of the classification, not clinical, not molecular |
| Autosomal dominant tubule interstitial kidney disease (67) | Meeting, agreement on the manuscript | Agreement on a novel name: ADTKD Classification based on the underlying genetic defect: ADTKD-gene | Provide information on the disease | Use in communication with patients may not be easy |
| Endocrine diseases (68) | Literature review | Groups of diseases by organ | | |
| Diabetes mellitus (MODY) (70) | Meetings, agreement on the manuscript | Groups of diabetes by diseases' mechanism | Provide information on the disease mechanism Allow numbering of new diabetes after identification of new genes for MODY (MODY1, MODY2, MODY3, MODY4, ...) | Very large groups of disease (type 2 diabetes for example) |
| Osteogenesis Imperfecta (64) | Literature review | Phenotypes on evolution, radiology, clinics and genetics | Provide information on the disease mechanism and genetics | Confusing as one causing gene may be in different categories |

Table 3: Definition of major and minor criteria for iPPSD and differential diagnoses


| | | Assessment | Differential diagnosis | References |
|---------------------------------|---|---|--|-------------------------------|
| I. MAJOR CRITERIA | 1. PTH resistance | Ionized calcium, total calcium Phosphate Magnesium PTH Vitamin D (25OHD) Creatinine Urinary calcium Urinary phosphate PTH infusion test in challenging cases | Normocalcemic hyperparathyroidism Renal failure Vitamin D deficiency or any kind of secondary hyperparathyroidism | (16) |
| | 2. Ectopic ossification | Detailed physical exam X-rays | Fibrodysplasia Ossificans Progressiva (FOP, OMIM# 135100), Post-traumatic Osteoma Cutis | |
| | 3. Brachydactyly type E (comprises the IV) | Clinical inspection (fist), Hand and feet X-rays | Turner syndrome, Tricho-rhino-phalangeal syndrome (TRPS), TRPS I, (OMIM#190350), TRPS-II (OMIM#150230) and TRPS-III, (OMIM#190351) | |
| II. MINOR CRITERIA | 1. TSH resistance | TSH, T4I, Antibodies, Imaging ¹ | Mutations in the TSH receptor | (26,27) |
| | 2. Other hormonal resistances | IGF-1 (GH stimulation test if necessary), Calcitonin, LH, FSH, GnRH test | | (2,27,78,98–101) |
| | 3. Motor and cognitive retardation or impairment | Computed tomography scan and/or MRI of the brain, Psychopathological rating scales adjusted for age | | (24,25,34,85, 86,102,116,117) |
| | 4. Intrauterine and postnatal growth retardation | IUGR: Gestational age, birth weight, birth length, head circumference, comparison to reference charts; Post-natal growth: Growth charts, X-ray of the left hand for determination of the bone age | | (16,40,92,103,104,121) |
| | 5. Obesity/overweight | Weight SDS, BMI percentile, BMI z-score | | (23,105,106) |
| | 6. Flat nasal bridge and/or maxillar hypoplasia and/or round face | Clinical inspection | | (4,84,86,90) |
| iPPSD CLINICAL DIAGNOSIS | a) Presence of one major criteria, either number 1 or 2; b) Presence of major criteria number 3 and at least 2 minor criteria ² | | | |

¹ US in adults with hypothyroidism and no evidence for autoimmunity; thyroid imaging through thyroid scintigraphy and US in neonates diagnosed through screening for congenital hypothyroidism.

² Minor criteria are nonspecific (obesity/cognitive impairment); for instance, the association of BDE + obesity or BDE + cognitive impairment would not be relevant for our classification. By raising the number of minor criteria from 1 to 2, we will reduce the risk of overdiagnosing patients with iPPSD.

PPSD

PTH/PTHrP






Figure 2




Figure 3




Figure 4